

MAP EXPLANATION

This map depicts a direct reclassification of the surficial geologic map of this area (House and others, 2006) into a series of relative flood hazard classes. The flood hazard classes are defined and are not intended to represent or support administrative or regulatory flood zone boundaries. They do not have specific implications for flood insurance. The hazard zones depict the local and spatial variability of flood hazard zones as inferred from geologic evidence. The hazard zones are based on surficial geologic stability and geomorphic position that form the basis of the geologic map. The classification scheme generally represents a conservative interpretation of the geologic data.

VERY HIGH Areas of the most frequent and concentrated runoff or erosion. Includes well-defined active channels, broad gravelly, and sparsely vegetated zones of diffuse distributary flow networks on active alluvial fans and high-velocity sheetflow on piedmont channels and playas (shown with a stippled pattern). Processes include high-velocity, channelized flow and high-velocity sheetflow on piedmont channels and playas. Channel boundaries and positions are generally unstable and may shift considerably by standing water on a regular basis. Piedmont areas are not subject to high-velocity flows, but do experience extensive channelization and erosion by standing water on a regular basis. Piedmont areas are subject to flooding from the toes of adjacent active alluvial fans and channels. Processes of sediment erosion, transport, and deposition in these areas are vigorous and involve particle sizes ranging from coarse gravel and cobbles to sand and silt. Corresponding surficial morphology includes prominent alluvial channels, fresh gravel bars, and relatively flat gravel sheets and coarse sand and silt. Corresponding surficial morphology includes prominent alluvial channels, fresh gravel bars, and relatively flat gravel sheets and coarse sand and silt. Corresponding surficial morphology includes prominent alluvial channels, fresh gravel bars, and relatively flat gravel sheets and coarse sand and silt. This class is Modern to late Holocene in age (0 to approximately 4000 years).

HIGH Areas of frequent, concentrated to widespread, relatively unchanneled runoff. Commonly includes well-defined active channels, broad gravelly, and sparsely vegetated zones of diffuse distributary flow networks on active alluvial fans, low-lying channel-bounding terraces, and intermediately active alluvial fan areas. Low-lying channel-bounding terraces and intermediately active alluvial fan areas are generally stable and are not subject to overflow and reoccupation by active channel networks. Sediment characteristics similar to areas in the very high hazard class. These areas have a high potential to convey flow during large floods because of their generally high relief, which precludes a lower hazard determination. Channel and flow network boundaries are generally stable. Morphology characteristics include relatively fresh bar and channel to slightly weathered bar and swale complexes, relatively fresh and stable alluvial fan surfaces, and relatively flat gravel sheets in playa fringe areas. Gravel and sand are locally in low-lying swales. Light rock varnish may be present on the surface of gravel. Geologic deposits in this class range from Modern to late Holocene (0 to approximately 4000 years). Geologic evidence indicates this class has high potential to convey dangerous flows during large flood events. It poses a significant floodplain management concern.

MODERATE Areas of intermediately mixed, highly active alluvial surfaces, intermediately active or recently abandoned alluvial surfaces, and dispersed remnants of stable alluvial fans. This class includes active and recently abandoned (last 100 to few 1000s of years) alluvial surfaces, stable alluvial fans, and stable alluvial fan surfaces. These surfaces are generally stable and are not subject to overflow and reoccupation by active channel networks. Sediment characteristics include weathered bar and swale complexes with muted characteristics similar to areas in the high hazard class. Morphology characteristics include weathered bar and swale complexes with muted characteristics similar to areas in the high hazard class. Morphology characteristics include weathered bar and swale complexes with muted characteristics similar to areas in the high hazard class. This class also includes broad areas of planar alluvial fan surfaces marked with at least 0.75 m of windblown sand which is commonly overlain by a thin and loose gravel lag. The preponderance of silt and clay in these surfaces is particularly widespread in Hidden Valley and along the north and east sides of Jean and Roach Lakes. Soil development on the sedimented surfaces is minimal and characterized by weak cambic (Bv) and calcic (Ck) soil horizons. Surface morphology, soil development, and relations to regional studies strongly suggest that surfaces in this class have been subject to alluvial fan processes for at least the last 7000 to 8000 years, and flood hazards are not significant except locally where remnants of older, active channels, in these areas, particularly in the Ivanpah Valley and along the north and east sides of Jean and Roach Lakes, may be present. Members of this class exhibit strongly planar surfaces with moderately to strongly eroded, channelized, and locally planar, relatively unchanneled surfaces. Surface clasts of carbonate rocks are weakly to moderately etched, channelized, and locally planar. Soil development associated with these surfaces is characterized by strong Av and Bw horizons and slope 1 to 2% to horizon development. This class represents a moderate floodplain management concern.

LOW Areas of stable alluvial surfaces that have been largely eroded from active alluvial fan processes for more than 5000 years. Members of this class, however, are linked too closely in space and time with areas in the high and moderate classes to assign them to flood hazard zones. Members of this class exhibit strongly planar surfaces with moderately to strongly eroded, channelized, and locally planar, relatively unchanneled surfaces. Surface clasts of carbonate rocks are weakly to moderately etched, channelized, and locally planar. Soil development associated with these surfaces is characterized by strong Av and Bw horizons and slope 1 to 2% to horizon development. This class also includes broad areas of planar alluvial fan surfaces marked with at least 0.75 m of windblown sand which is commonly overlain by a thin and loose gravel lag. The preponderance of silt and clay in these surfaces is particularly widespread in Hidden Valley and along the north and east sides of Jean and Roach Lakes. Soil development on the sedimented surfaces is minimal and characterized by weak cambic (Bv) and calcic (Ck) soil horizons. Surface morphology, soil development, and relations to regional studies strongly suggest that surfaces in this class have been subject to alluvial fan processes for at least the last 7000 to 8000 years, and flood hazards are not significant except locally where remnants of older, active channels, in these areas, particularly in the Ivanpah Valley and along the north and east sides of Jean and Roach Lakes, may be present. Members of this class exhibit strongly planar surfaces with moderately to strongly eroded, channelized, and locally planar, relatively unchanneled surfaces. Surface clasts of carbonate rocks are weakly to moderately etched, channelized, and locally planar. Soil development associated with these surfaces is characterized by strong Av and Bw horizons and slope 1 to 2% to horizon development. This class represents a lower floodplain management concern than preceding ones. It is locally vulnerable to overflow and lateral accretion.

VERY LOW This class includes geologic deposits and surfaces that do not experience alluvial flooding, but may be subject to flooding from local sources. It includes thick, actively accumulating mantles of eolian sand and silt, moderately to very strongly carbonate-cemented, relict alluvial surfaces that have been free from active alluvial fan processes for periods well in excess of thousands of years (House and others, 2006). In some cases, surfaces included in this class are so high-standing as to obviously preclude alluvial fan flood hazards. In other cases, low-standing members exhibit surface morphology and soil development characteristics that are consistent with existing long periods of stability, including planar surface remnants with light, darkly varnished gravel pavements that may include abundant clasts of coarse silt, moderately to deeply weathered surface clasts (cliffing, pitting, and spalling), deeply buried planar to weakly convex surface remnants with eroded calcic soil horizons, high-standing, deeply incised fan remnants with retrograde gravel pavements on surface and sub-surface, and high-standing planar surfaces underlain by massive petrocalcic soils. Eolian features included in this class include locally thick accumulations of sand on ancient fan and bedrock surfaces, rock sand ramps that overlie steep mountain-front slopes, and some small areas of dunes. This class does not represent a significant floodplain management problem, but some local conditions may merit concern.


VARIABLE Small areas that may have special hazardous conditions that are not linked to alluvial fan flood hazards. This includes mixed cultural gravel and debris flow deposits on steeply sloping hillslopes and variably active talus piles and cultural debris cores below steep bedrock cliffs. The former situation is common on steep slopes on volcanic rocks of the McCullough Range, particularly in the Hidden Valley area, and below Table Mountain in the southern Spring Mountains. The latter situation is mainly present in parts of the Bird Spring Range and Spring Mountains where steep bedrock cliffs are common. These types of units mainly the latter type were mapped separately throughout the area where they were easily distinguished on aerial photographs and other imagery (House and others, 2006). Members of this class represent special situations and should be evaluated on an individual basis. They comprise a very small part of the map area and are often in rugged and remote settings.

INDERMINATE Areas that have been extensively modified by excavation, artificial fill, or commercial development. Flood hazards in these areas cannot be assessed from the basis of geologic evidence. This includes the course of Interstate 15, the Union Pacific railroad, borrow pits and mining operations, and developed areas in Goodsprings, Jean (including the Jean Airport and the correctional facility), and Primm. This class represents a special situation. It may locally represent a significant floodplain management concern, but large tracts of it are not flood zones.

UNMAPPED Areas mapped as bedrock by House and others (2006). Locally includes small areas of active and inactive alluvial surfaces, colluvium, and minor amounts of eolian sediment. Significant flood and debris flow hazards exist in narrow bedrock canyons and steep slopes in these areas but are too small to map. Extremely rugged topography in most bedrock areas limits potential for commercial and residential development. This class may locally represent a floodplain management concern, but very large tracts of it are neither flood-prone nor readily accessible.

REFERENCES
House, P.K., 2005. Using geology to improve flood hazard management on alluvial fans: an example from Laughlin, Nevada. Proceedings of the American Water Resources Association, v. 41, no. 6, p. 1431-1447.
House, P.K., Rameck, A.S., and Brock, B.J., 2006. Surficial geologic map of the Ivanpah Valley area, Clark County, Nevada. Nevada Bureau of Mines and Geology Map 158, 1:50,000.

7.5' Quadrangle Index and Basin Location Map



Map boundary	Other road or street	Township and range lines
Interstate highway	Trail	State boundary
State highway	Railroad	Powerline

NEVADA BUREAU OF MINES AND GEOLOGY
MACKAY SCHOOL OF EARTH SCIENCES AND ENGINEERING
UNIVERSITY OF NEVADA, RENO
Funding from Clark County Regional Flood Control District and U.S. Geological Survey (STATEMAP Program, Agreement No. 04-HQ-A0-0032, 05-HQ-A0-0049, and 02-HQ-A0-0019). Field work performed from 2002-2006.
We thank the following individuals for their assistance and input towards the development of this map: Brian Park, Thomas Prescott, Amy Brock, and Andrew Hanson.
Chief Editor: John W. Hawley
Field review by: John W. Hawley (Hawley Geomatters, Albuquerque, NM); Philip A. Penland (USGS, John W. Bell (NSBG); Catherine P. Kyle House, Christine M. Antill and Elizabeth C. Couse (Clark County Regional Flood Control District); Douglas J. Menker (HRCS, Reno, NV); John W. Bell (NSBG); Douglas J. Menker (HRCS, Reno, NV); John W. Bell (NSBG); Douglas J. Menker (HRCS, Reno, NV); John W. Bell (NSBG); Douglas J. Menker (HRCS, Reno, NV).
First Edition, 2007
Revised Edition, 2007
Copyright © 2007 by Nevada Bureau of Mines and Geology
University of Nevada, Mail Stop 1775
Reno, Nevada 89512-1775
(775) 784-6691
http://www.nbgm.edu; www.nvbgm.edu

GEOLOGIC ASSESSMENT OF PIEDMONT AND PLAYA FLOOD HAZARDS IN THE IVANPAH VALLEY AREA, CLARK COUNTY, NEVADA

P. Kyle House
2007

