

Petroleum Data Map of Nevada

Larry J. Garside and Ronald H. Hess
Nevada Bureau of Mines and Geology
2007

OIL AND GAS WELL DATA

Current and past oil production wells, petroleum exploration wells with shows of oil and/or gas, and petroleum exploration wells with no shows or no data (from Hess, 2004), updated through 2006, are plotted on the map.

SEEPS

The term "seep" is used to refer to shows of oil, gas, or solid bitumen at the surface, in springs, and in non-petroleum wells. Data are from Garside and others (1988), and Barker (1996), Hulen and others (1994, 1999), and Schalla and others (1994).

SOURCE ROCKS

A number of stratigraphic units in Nevada have been mentioned as potential source rocks for petroleum. We list below, by age, these formations, the map unit descriptors of Crafford (2007) for the map units that represent or include the potential source rocks, and select references that mention or discuss source-rock potential of the referenced units. Some of the units of Crafford (2007) are more inclusive than the formations listed below. The outcrops of these units from Crafford (2007) are shown collectively on the map.

Miocene(?) Lower Humboldt Formation of Fritch and Pekarek (1994) (outcrops in the Windermere Hills northeast of Wells) – T₁ as shown by Coats (1987). Reference: Fritch and Pekarek, 1994.

Eocene/Oligocene(?) Elko Formation – T₁. References: Poole and others, 1983; Poole and Claypool, 1984; Palmer, 1984.

Paleocene/Eocene/Cretaceous(?) Sheep Pass Formation – T₁. References: Poole and others, 1983; Poole and Claypool, 1984.

Late Cretaceous/Early Tertiary Pansy Lee Formation – TK₉ (Humboldt County only). Reference: Wilden, 1979.

Cretaceous Newark Canyon Formation – K₉ (in Eureka, White Pine, Elko, and Nye Counties). Reference: Poole and others, 1983.

Cretaceous King Lear Formation – K₉ (Humboldt County only). Reference: Wilden, 1979.

Triassic Favret Formation (part of the Star Peak Group) – TR₁. Reference: Cook, 1987.

Permian Phosphite Formation (Park City Group) – P₁. Carbonaceous shale members in eastern Elko County as shown by Maughan, 1984, fig. 2). References: Maughan, 1984; Poole and Claypool, 1984, p. 196; Wardlaw and others, 1979.

Pennsylvanian/Permian Bird Springs Formation – PIP₁ (Clark County only). Reference: Longwell and others, 1965, p. 160.

Mississippian Chatham Shale, Webb Formation, Woodruff Formation – IP₁MD₁. Reference: Poole and others, 1983; Poole and Claypool, 1984.

Devonian/Mississippian Pilot Shale – PIP₁, MD₁, and some MD₂. Reference: Poole and Claypool, 1984, p. 196.

Ordovician Vintni Formation (?) – DO₁. References: Poole and others, 1983; Poole and Claypool, 1984.

CONDONOT COLOR ALTERATION INDICES (CAI)

Condonots are the apatitic hard parts (teeth) of an extinct group of marine worm-like animals. They range in age from Cambrian to Triassic and are excellent biostratigraphic zone indicators. They contain trace amounts of organic matter that change color with increasing temperature. Condonot color alteration indices (CAI), reported in increments of 0.5, are a measure of organic and mineral metamorphism (Epstein and others, 1977). Thus, these indices are useful in interpreting the heating history of petroleum source rocks. Because thermal history is a key factor in hydrocarbon generation and preservation, CAIs (in combination with source-rock organic content) can be useful in evaluating petroleum potential. Harris and Crafford (2007) reported maximum and minimum CAI values for collections of Nevada condonots. Most collections did not exhibit a range, and the maximum and minimum values were the same. However, some samples exhibited a range of 0.5 to 4 or more. We plotted the maximum values reported by Harris and Crafford (2007). We suggest that if any of the condonots were heated to a higher level in the host rock, then any associated hydrocarbons would have also reached those temperatures (e.g., Repetti and others, 2005). In general, CAI > 4.5 is considered to be the thermal cutoff for most hydrocarbon production (Harris and others, 1980).

The ages of the condonots (and thus the rocks sampled) were represented using the AgeMin numerical column in Harris and Crafford (2007): <3099 – Triassic; 3100–3299 – Pennsylvanian/Permian; 3300–3499 – Mississippian/Devonian; >3500 – pre-Devonian.

REFERENCES

Barker, C.E., 1996, Resource assessment of the Bureau of Land Management's Winnemucca District and Surprise Resource Area, Northwest Nevada and Northeast California—geochemical analysis and thermotectonic modeling to evaluate conceptual petroleum plays. U.S. Geological Survey Open-File Report 96-051, 51 p.

Coats, R.R., 1987, Geology of Elko County, Nevada. Nevada Bureau of Mines and Geology Bulletin 101, 112 p.

Cook, H.E., 1987, National Petroleum Assessment, western Basin and Range province. U.S. Geological Survey Open-File Report 87-450L.

Crafford, A.E.J., 2007, Geologic map of Nevada. U.S. Geological Survey Data Series DS-249.

Epstein, A.G., Epstein, J.B., and Harris, L.D., 1977, Condonot color alteration—an index to organic metamorphism. U.S. Geological Survey Professional Paper 995, 27 p.

Fritch, W.E., and Pekarek, A.H., 1994, Lower Miocene petroleum potential of northeast Elko County, Nevada. In Schalla, R.A., and Johnson, E.H., eds., Oil fields of the Great Basin. Nevada Petroleum Society, Reno, p. 151-162.

Garside, L.J., Hess, R.H., Fleming, K.L., and Weimer, B.S., 1988, Oil and gas developments in Nevada. Nevada Bureau of Mines and Geology Bulletin 104, 136 p. www.nbmg.unr.edu/ol4b104.htm

Harris, A.G., and Crafford, A.E.J., 2007, A conodont database of Nevada. In Crafford, A.E.J., Geologic map of Nevada. U.S. Geological Survey Data Series DS-249.

Harris, A.G., Wardlaw, B.R., Rust, C.C., and Merrill, G.K., 1980, Maps for assessing thermal maturity (condonot color alteration index maps) in Ordovician through Triassic rocks in Nevada and Utah and adjacent parts of Idaho and California. U.S. Geological Survey Miscellaneous Investigation Series Map I-1249.

Hess, R.H., 2004, Nevada Oil and Gas Well Database (NVOGWD). Nevada Bureau of Mines and Geology Open-File Report 04-1242. www.nbmg.unr.edu/ol041242.htm with well information updated through 2006 for this map.

Hulen, J.B., Pinnel, M.L., Nelson, D.J., Cox, J.W., and Blake, J., 1994, The Yankee gold mine occurrence, Alligator Ridge District, Nevada: an enhanced and updated, petrogenetic reservoir. In Schalla, R.A., and Johnson, E.H., eds., Oil fields of the Great Basin. Nevada Petroleum Society, Reno, p. 131-142.

Hulen, J.B., Collier, J.W., Johnson, S.D., and Allen, R., 1999, Oils in the Dixie Valley and Kyle Hill Springs geothermal systems, Nevada—potentially sensitive indicators of natural and induced reservoir processes. Proceedings, Twenty-fourth Workshop on Geothermal Reservoir Engineering, Stanford University, January 25-27, 1999, 19 p. [GPR-TR-182].

Longwell, C.R., Pampayan, E.H., Bowyer, B., and Roberts, R.J., 1966, Geology and mineral deposits of Clark County, Nevada. Nevada Bureau of Mines and Geology Bulletin 62, 216 p.

Maughan, E.K., 1984, Geological setting and some geochemistry of petroleum source rocks in the Permian Phosphite Formation. In Woodward, J., Meissner, F.F., and Clayton, J.L., eds., Hydrocarbon Source Rocks of the Greater Rocky Mountain Region. Rocky Mountain Association of Geologists, Denver, Colorado, p. 281-294.

Palmer, S.E., 1984, Hydrocarbon source potential of organic facies of the lacustrine Elko Formation (Eocene/Oligocene), northeast Nevada. In Woodward, J., Meissner, F.F., and Clayton, J.L., eds., Hydrocarbon Source Rocks of the Greater Rocky Mountain Region. Rocky Mountain Association of Geologists, Denver, Colorado, p. 497-511.

Poole, F.G., and Claypool, G.E., 1984, Petroleum source-rock potential and crustal correlation in the Great Basin. In Woodward, J., Meissner, F.F., and Clayton, J.L., eds., Hydrocarbon Source Rocks of the Greater Rocky Mountain Region. Rocky Mountain Association of Geologists, Denver, Colorado, p. 277-283.

Poole, F.G., Claypool, G.E., and Fouch, T.D., 1983, Major episodes of petroleum generation in part of the northern Great Basin. Geothermal Resources Council Special Report 13, p. 207-213.

Repetti, J.E., Ryker, R.T., Avery, K.L., and Trippl, M.H., 2005, Thermal maturity patterns (CAI and %Ro) in the Ordovician and Devonian rocks of the Appalachian basin in West Virginia. U.S. Geological Survey Open-File Report 2005-1078, 89 p.

Schalla, R.A., Barker, C.E., and Neumann, W.H., 1994, Hot water and oil from a shallow exploratory borehole, Basin and Range Province, Nevada. In Schalla, R.A., and Johnson, E.H., eds., Oil fields of the Great Basin. Nevada Petroleum Society, Reno, p. 127-130.

Wardlaw, B.R., Collinson, J.W., and Ketter, K. B., 1979, Regional relations of middle Permian rocks in Idaho, Nevada, and Utah. In Newman, G.W., and Goode, H.E., eds., Basin and Range Symposium. Rocky Mountain Association of Geologists and Utah Geological Association, p. 277-283.

Wilden, Ronald, 1979, Petroleum exploration possibilities in northwestern Nevada. In Newman, G.W., and Goode, H.E., eds., Basin and Range Symposium. Rocky Mountain Association of Geologists and Utah Geological Association, p. 541-548.

Oil and Gas Wells through 2006

- Oil producer
- Oil and gas show
- Oil show
- Gas show
- No show or no data

Seeps and Other Shows

- Gas seep
- Gas show in water well or spring
- Oil seep or hydrocarbon occurrence
- Oil show in water well or spring

Source Rock Outcrops (from Crafford, 2007)

CAI Maximum and Age

Age	CAI Maximum				
	CAI 1.0 & 1.5	CAI 2.0 & 2.5	CAI 3.0 & 3.5	CAI 4.0 & 4.5	CAI > 4.5
Triassic	●	●	●	●	●
Pennsylvanian-Permian	▲	▲	▲	▲	▲
Mississippian-Devonian	■	■	■	■	■
pre-Devonian	▼	▼	▼	▼	▼

Nevada Bureau of Mines and Geology
Mackay School of Earth Sciences and Engineering
College of Science
University of Nevada, Reno


Reviewed by William J. Ehn, Charles W. Gillespie, Jonathan G. Price, and Jerry P. Walker
Edited by Dick Meeuwig

First Edition, 2007
Printed by Nevada Bureau of Mines and Geology

This map was printed on an electronic order delivery from digital files. Limited color calibration may vary between electronic prints and a set of 500 copies of the same plates, and paper map change rate due to electronic printing, color and quantities may not be exact on paper maps.

Digital materials and pdf copy of this map available at www.nbmg.unr.edu/ol04

For sale by:
Nevada Bureau of Mines and Geology
University of Nevada, Reno 178
Reno, Nevada 89557-0178
(775) 784-4901, ext. 2
www.nbmg.unr.edu; nbmg@unr.edu



Scale 1:1,000,000
1 inch equals approximately 16 miles
Base Map: Modified from NBMG Map 43, 1995