

630 000 FEET

36°07'30"

**NEVADA BUREAU OF MINES AND GEOLOGY** 

LAS VEGAS AREA MAP 3Bg





Qa Alluvium of active washes Pink to pale-brown fine sand to pebble to cobble gravel occurring in incised, active stream solidated to locally cemented by petrocalcic cement (case hardened). Clasts are mostly limestone and dolomite in the northern and central parts of the quadrangle and volcanic rock fragments in the southern part. Sand-size sediment is mainly fragmental limestone and dolomite; quartz and feldspar are subordinate and detrital gypsum locally is an important compo-

Sheetwash alluvium Pink to brown sand, pebble to cobble gravel, and petrocalcic fragments occurring as thin veneers downslope from fault scarps; unconsolidated to slightly consolidated. Recycled from rock units exposed on upthrown sides of fault scarps. Active; subject to reworking by wind. Sand dunes Pink to pale-brown fine to medium sand occurring in small to large barchan dunes and dune fields; unconsolidated to moderately consolidated. Intermittently active to inactive and stabilized deposits derived from eolian reworking of Qa, Qai, Qoa, and Qt. Composition is predominantly detrital limestone, dolomite, and gypsum with subordinate

quartz and feldspar. Intermittently active alluvium Pink to pale-brown sand and pebble to cobble gravel occurring mainly on between-channel alluvial flats and less commonly in incised washes; slightly to moderately consolidated. Deposits are lithologically similar to Qa; carbonate clasts are predominant in the northern and central parts of the quadrangle and volcanic clasts are predominant in the southern part. Unit grades laterally into Qa; contact between Qai and Qa in some places is arbitrary and may shift. Contains sediment transported and deposited during highwater stages or heavy flood discharges.

Older alluvium of Blue Diamond and Red Rock fans Pink to brown pebble to small cobble gravel with subordinate pebblebearing sand; moderately to well consolidated to locally cemented; petrocalcic carbonate horizon (calcrete) 1 ½ - 2 m thick occurs at or near the surface. Clasts are prelimestone and dolomite with subordinate quartzite. Surface of unit is slightly to well dissected, lacks channel-and-bar morphology, has well-developed desert pavement, and is dark colored and mottled on aerial photographs. Surface clasts have slight to moderate desert varnish; carbonate clasts are typically etched. Unit thickens westward to exposed thicknesses in excess of 5 m; unit feathers out to the east on upthrown fault

blocks. Stippled where degraded and calcrete absent or weakly developed. Deposits of Duck Creek Terrace Pink to pale-brown pebble to cobble gravel, pebbly sand, and subordinate fine to coarse sand; moderately to well consolidated. Clasts include limestone, dolomite, quartzite, and volcanic rock fragments in varying proportions. Deposits occur in terraces along Paradise Valley and are related to backfilling and wash incision along Duck Creek.

Qog, Older gravel deposits Dissected, consolidated to strongly Qog<sub>2</sub> cemented gravel deposits that represent early-generation Qog<sub>3</sub> alluvial-fan development derived from outside the quadrangle; within the quadrangle, the deposits feather out on top of QTs. Qog1: Light-gray to pink-brown, imbricated, clast-supported pebble to cobble gravel capped by a matrix-supported petrocalcic (calcrete) horizon more than 3 m thick. Clasts mostly consist of limestone and dolomite with subordinate quartzite. At the surface, carbonate clasts are strongly etched and broken and possess slight to moderate petrocalcic coatings; the petrocalcic horizon is eroded and petrocalcic fragments form varying proportions of the surface clasts. Unit feathers out to the east and thickens westward to exposed thicknesses greater than 10 m. Qog2: Light- to medium-brown sandy pebble to cobble gravel capped by a petrocalcic

(calcrete) horizon more than 2 m thick. Clasts are predominantly basalt and andesite with subordinate limestone, dolomite, and quartzite. The petrocalcic horizon occurs at the surface except in the extreme southeast part of the quadrangle where it is covered by a thin (1-2 m) veneer of bouldery alluvium (indicated by stippled pattern). Surface of unit has moderately to well-developed desert varnish; carbonate clasts are etched; bouldery veneer exhibits subdued to moderately preserved channel-and-bar morphology. Qog: Light- to medium-brown pebble to boulder gravel interstratified with pebbly fine calcareous sand; capped by a petrocalcic (calcrete) horizon more than 2 m thick. Gravel clasts are predominantly limestone and dolomite pebbles and cobbles. Surface of deposit is rounded and mantled with etched and broken car-


bonate fragments. QTs Consolidated sediments White and light-gray to light-and pale-red fine sand interstratified with silt, pebbly sand, and pebble to small cobble gravel; moderately to well consolidated to strongly cemented. Layers of petrocalcic carbonate are common and have variable textures and fabrics; surface exposures are locally capped by a resistant petrocalcic crust. Fibrous and encrusting gypsum are common. Massive to flat laminated to shallowly trough laminated; scour-and-fill channel structures occur locally. Unit was deposited by streams; lacustrine sediments have not been documented.

Contact Dashed where approximately located. Hachures indicate that a geologic unit occurs in an incised channel or in a cut-and-fill terrace or that a younger geologic unit truncates an older geologic unit with but-

tress unconformity.

- Fault scarp Ball on downdropped side of escarpment Urbanized area or artificial cut and/or fill for which geologic data are not available

Jonathan C. Matti and Fred W. Bachhuber, 1985 Supplementary mapping was provided by John W. Bell



First edition, first printing, 1985: 1000 copies
Printing: Williams and Heintz Map Corp., Washington, D.C.
Editing: Bridgett Boulton and Alice Sjoberg
Cartography: Larry Jacox
Cofor separation assistance: Michael Tracy
Typesetting: Rayetita Buckley
Pasteup: Mati Stephens For sale by the Nevada Bureau of Mines and Geology, University of Nevada-Reno, Reno, Nevada, 89557-0088 Order Map 3Bg, \$4.00